

# Pemetrexed: Alimta®; Pemfexy™; Pemrydi RTU™; Pemetrexed Ψ (Intravenous)

-E-

Document Number: MODA-0489

**Last Review Date: 07/02/2024**

**Date of Origin: 08/05/2019**

**Dates Reviewed: 08/2019, 10/2019, 01/2020, 04/2020, 07/2020, 10/2020, 01/2021, 05/2021, 07/2021, 10/2021, 02/2022, 04/2022, 07/2022, 10/2022, 01/2023, 04/2023, 08/2023, 10/2023, 01/2024, 04/2024, 07/2024**

## I. Length of Authorization <sup>15,26,28-30</sup>

Coverage will be provided for 6 months and may be renewed, unless otherwise specified.

- Thymomas: Coverage will be provided for six (6) 21-day cycles and may NOT be renewed.
- Mesothelioma (PM) in combination with bevacizumab AND either cisplatin or carboplatin: Coverage will be provided for six (6) cycles and may NOT be renewed.

## II. Dosing Limits

### A. Quantity Limit (max daily dose) [NDC Unit]:

- Alimta 100 mg powder for injection in a single-use vial: 4 vials every 21 days
- Alimta 500 mg powder for injection in a single-use vial: 4 vials every 21 days
- Pemfexy 500 mg solution for injection in a multi-dose vial: 4 vials every 21 days
- Pemetrexed 750 mg powder for injection: 2 vials every 21 days
- Pemetrexed 1000 mg powder for injection: 2 vials every 21 days
- Pemetrexed 100 mg/4 mL solution for injection: 4 vials every 21 days
- Pemetrexed 500 mg/20 mL solution for injection: 4 vials every 21 days
- Pemetrexed 850 mg/34 mL solution for injection: 2 vials every 21 days
- Pemetrexed 1000 mg/40 mL solution for injection: 2 vials every 21 days
- Pemrydi RTU 100 mg/10 mL solution for injection: 4 vials every 21 days
- Pemrydi RTU 500 mg/50 mL solution for injection: 4 vials every 21 days
- Pemrydi RTU 1000 mg/100 mL solution for injection: 2 vials every 21 days

### B. Max Units (per dose and over time) [HCPCS Unit]:

- Pemfexy (500 mg MDV):
  - Ovarian Cancer: 225 billable units every 21 days
  - Thymomas: 125 billable units every 21 days
  - All other indications: 125 billable units every 21 days

- Pemetrexed (all other manufacturers) (100 mg, 500 mg, 750 mg, 850 mg, and 1000 mg SDV):
  - Ovarian Cancer: 230 billable units every 21 days
  - Thymomas: 130 billable units every 21 days
  - All other indications: 130 billable units every 21 days

### III. Initial Approval Criteria <sup>1-3</sup>

Coverage is provided in the following conditions:

- Patient must have a contraindication, intolerance, or failure to ALL alternative pemetrexed products prior to consideration of Pemfexy (J9304); **AND**

- Patient is at least 18 years of age; **AND**

#### Peritoneal Mesothelioma (PeM) ‡ <sup>4,30</sup>

- Used in combination with cisplatin or carboplatin (if cisplatin ineligible); **AND**
  - Used as adjuvant therapy; **AND**
    - Patient has unicavitary disease with epithelioid histology; **AND**
    - Patient has surgical/pathologic high-risk features\*\* and no neoadjuvant therapy was given; **OR**
  - Used as first-line therapy; **AND**
    - Patient has biphasic/sarcomatoid histology or bicavitary disease; **OR**
    - Patient has unicavitary disease with epithelioid histology; **AND**
      - Patient is medically inoperable and/or complete cytoreduction is not achievable (including high-risk features\*\*); **OR**
      - Patient has recurrent disease after prior cytoreductive surgery (CRS) + hyperthermic intraperitoneal (IP) chemotherapy (HIPEC) and no previous adjuvant systemic therapy was given; **OR**
  - Used as subsequent therapy; **AND**
    - Immunotherapy (i.e., nivolumab/ipilimumab) was administered as first-line treatment

\*\* High-risk features include Ki-67 >9%, nodal metastasis, high tumor burden (Peritoneal Cancer Index [PCI] >17), completeness of cytoreduction (CC) score >1, biphasic disease, or bicavitary disease

#### Pleural Mesothelioma (PM) † ‡ Φ <sup>1-7,11,27,79e,80e</sup>

- Used as induction therapy; **AND**
  - Used in combination with cisplatin or carboplatin (if cisplatin ineligible) in patients with clinical stage I-IIIa disease and epithelioid histology; **OR**
- Used as first-line therapy; **AND**

- Used in combination with bevacizumab AND either cisplatin or carboplatin (if cisplatin ineligible); **AND**
  - Patient has unresectable disease; **OR**
- Used in combination with cisplatin or carboplatin (if cisplatin ineligible); **AND**
  - Disease is unresectable or patient is not a candidate for curative surgery; **OR**
- Used as subsequent therapy; **AND**
  - Used as a single agent OR in combination with cisplatin or carboplatin (if cisplatin ineligible), with or without bevacizumab; **AND**
    - Immunotherapy (i.e., nivolumab/ipilimumab) was administered as first-line treatment; **OR**
    - Used as a rechallenge if pemetrexed-based treatment was administered first-line with good response

**Non-Squamous Non-Small Cell Lung Cancer (NS-NSCLC) † ‡** <sup>1-4,8-10,12,13,23,29,31,50e,51e,54e,56e-58e,81e-83e,92e-95e,98e</sup>

- Used in combination with carboplatin or cisplatin; **OR**
- Used in combination with bevacizumab, pembrolizumab, cemiplimab, or durvalumab for continuation maintenance therapy if previously used first-line and patient achieved a tumor response or stable disease following initial therapy; **OR**
- Used in combination with nivolumab and either cisplatin or carboplatin as neoadjuvant therapy for resectable (tumors  $\geq$  4 cm or node positive) disease; **OR**
- Used in combination with pembrolizumab and cisplatin as neoadjuvant therapy for resectable (tumors  $\geq$  4 cm or node positive) disease; **OR**
- Patient has recurrent, advanced, or metastatic disease (excluding locoregional recurrence or symptomatic local disease without evidence of disseminated disease) or mediastinal lymph node recurrence with prior radiation therapy; **AND**
  - Used in combination with cemiplimab and either cisplatin or carboplatin; **OR**
  - Used in combination with osimertinib as first-line therapy for EGFR exon 19 deletion or exon 21 L858R mutation positive disease; **OR**
  - Used in combination with amivantamab-vmjw and carboplatin as first-line therapy for EGFR exon 20 insertion mutation positive disease; **OR**
  - Used in combination with amivantamab-vmjw and carboplatin following disease progression on osimertinib for EGFR exon 19 deletion or exon 21 L858R mutation positive disease; **OR**
  - Used in combination with pembrolizumab and either cisplatin or carboplatin; **OR**
  - Used in combination with tremelimumab, durvalumab, and either cisplatin or carboplatin; **OR**
  - Used in combination with nivolumab, ipilimumab, and either cisplatin or carboplatin; **OR**

- Used as a single agent; **AND**
  - Used as first-line therapy for tumors that are negative for actionable molecular biomarkers\* †; **OR**
  - Used as first-line therapy for EGFR exon 20 mutation, BRAF V600E-mutation, NTRK1/2/3 gene fusion, MET exon 14 skipping mutation, RET rearrangement, or ERBB2 (HER2) mutation positive tumors; **OR**
  - Used as subsequent therapy; **OR**
  - Used as continuation or switch maintenance therapy in patients who have achieved a tumor response or stable disease following initial platinum-based therapy

*\* Note: Actionable molecular genomic biomarkers include EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, and ERBB2 (HER2). Complete genotyping for EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, and ERBB2 (HER2), via biopsy and/or plasma testing. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. Treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.*

† May also be used for patients with KRAS G12C mutation positive tumors.

#### **Thymomas ‡** <sup>4,15,16,26,68e</sup>

- Used as a single agent; **AND**
- Used as second-line therapy; **AND**
- Patient has unresectable or metastatic disease

#### **Ovarian, Fallopian Tube, and Primary Peritoneal Cancer ‡** <sup>4,14,25,74e,75e</sup>

- Used as a single agent; **AND**
- Patient has platinum-resistant disease; **AND**
  - Patient has recurrent or persistent Grade 1 Endometrioid Carcinoma, Carcinosarcoma (Malignant Mixed Müllerian Tumors), Mucinous Carcinoma of the Ovary, Epithelial Ovarian/Fallopian Tube/Primary Peritoneal Cancer, or Clear Cell Carcinoma of the Ovary; **AND**
    - Patient is not experiencing an immediate biochemical relapse (i.e., rising CA-125 without radiographic evidence of disease); **OR**
  - Patient has recurrent Low-Grade Serous Carcinoma

**Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.**

## **§ Genomic Aberration/Mutational Driver Targeted Therapies**

| <b>(Note: not all inclusive, refer to guidelines for appropriate use)</b>                                    |   |   |  |
|--|---|---|--|
| <i>EGFR</i> exon 19 deletion or exon 21 L858R tumors   | <i>EGFR</i> S768I, L861Q, and/or G719X mutation positive tumors                             | <i>EGFR</i> exon 20 insertion mutation positive tumors                    | <i>NTRK1/2/3</i> gene fusion positive tumors                     |
| – Afatinib<br>– Erlotinib<br>– Dacomitinib<br>– Gefitinib<br>– Osimertinib<br>– Amivantamab                  | – Afatinib<br>– Erlotinib<br>– Dacomitinib<br>– Gefitinib<br>– Osimertinib<br>– Amivantamab | – Amivantamab   | – Larotrectinib<br>– Entrectinib                                 |
| <i>ALK</i> rearrangement-positive tumors   | <i>ROS1</i> rearrangement-positive tumors   | <i>BRAF</i> V600E-mutation positive tumors                                | <i>ERBB2 (HER2)</i> mutation positive tumors                     |
| – Alectinib<br>– Brigatinib<br>– Ceritinib<br>– Crizotinib<br>– Lorlatinib                                   | – Ceritinib<br>– Crizotinib<br>– Entrectinib<br>– Lorlatinib<br>– Repotrectinib             | – Dabrafenib ± trametinib<br>– Encorafenib + binimetinib<br>– Vemurafenib | – Fam-trastuzumab deruxtecan-nxki<br>– Ado-trastuzumab emtansine |
| PD-L1 tumor expression ≥ 1%  | <i>MET</i> exon-14 skipping mutations   | <i>RET</i> rearrangement-positive tumors                                  | <i>KRAS G12C</i> mutation positive tumors                        |
| – Pembrolizumab<br>– Atezolizumab<br>– Nivolumab + ipilimumab<br>– Cemiplimab<br>– Tremelimumab + durvalumab | – Capmatinib<br>– Crizotinib<br>– Tepotinib   | – Selpercatinib<br>– Cabozantinib<br>– Pralsetinib                        | – Sotorasib<br>– Adagrasib                                       |

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ◻ Orphan Drug

#### IV. Renewal Criteria <sup>1,2</sup>

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: myelosuppression (e.g., neutropenia, febrile neutropenia, thrombocytopenia, anemia), renal toxicity (CrCl < 45 mL/min), bullous and exfoliative skin toxicity (e.g., Stevens-Johnson Syndrome/Toxic epidermal necrolysis), interstitial pneumonitis, radiation recall, etc.; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**

#### Mesothelioma (PM) <sup>27,30</sup>

- Coverage may NOT be renewed when used in combination with bevacizumab AND either cisplatin or carboplatin

#### Thymomas <sup>16</sup>

- Coverage may NOT be renewed

## V. Dosage/Administration <sup>1-3,11,14,16,17,27,29-34</sup>

| Indication  | Dose  |
|---|---|
| Non-Squamous NSCLC  | Administer up to 500 mg/m <sup>2</sup> intravenously every 21 days  |
| Mesothelioma (peritoneal and pleural)   | Administer 500 mg/m <sup>2</sup> intravenously every 21 days <ul style="list-style-type: none"> <li>– For 6 cycles only when used in combination with bevacizumab AND either cisplatin or carboplatin</li> <li>– All others until disease progression or unacceptable toxicity</li> </ul> |
| Ovarian Cancer  | Administer 900 mg/m <sup>2</sup> intravenously every 21 days, until disease progression or unacceptable toxicity  |
| Thymomas  | Administer 500 mg/m <sup>2</sup> intravenously every 21 days for a maximum of 6 cycles or until disease progression or unacceptable toxicity  |
| <ul style="list-style-type: none"> <li>• Supplement with oral folic acid and intramuscular vitamin B12.</li> <li>• Avoid administration of ibuprofen for 2 days before, the day of, and 2 days following administration in patients with CrCl &lt;80 mL/min.</li> <li>• Do not administer in patients with CrCl &lt;45 mL/min.</li> </ul> |   |

## VI. Billing Code/Availability Information

| Product Formulation   | Drug                                     | Manufacturer | Type    | HCPCS Code    | NDC           |
|---|--|--------------|---------|---------------|---------------|
| Pemetrexed Disodium Hemipentahydrate Solution for injection | Pemrydi RTU 100 mg/10 mL SDV Ψ           | Lilly        | Brand   | J9324         | 70121-2453-xx |
|   | Pemrydi RTU 500 mg/50 mL SDV Ψ           |              |         |               | 70121-2461-xx |
|   | Pemrydi RTU 1000 mg/100 mL SDV Ψ         |              |         |               | 70121-2462-xx |
| Pemetrexed Disodium Lyophilisate for injection              | Alimta 100 mg powder for inj. SDV *      | Lilly        | Brand   | J9305         | 00002-7640-xx |
|   | Alimta 500 mg powder for inj. SDV *      |              |         |               | 00002-7623-xx |
|   | Pemetrexed 100 mg powder for inj. SDV Ψ  | Hospira      | Brand   | J9294         | 00409-1060-xx |
|   | Pemetrexed 500 mg powder for inj. SDV Ψ  |              |         |               | 00409-1061-xx |
|   | Pemetrexed 750 mg powder for inj. SDV *  | N/A          | Generic | J9305         | N/A           |
|   | Pemetrexed 1000 mg powder for inj. SDV * |              |         |               |               |
|   | Pemetrexed 100 mg powder for inj. SDV Ψ  | BluePoint    | Brand   | J9322         | 68001-0543-xx |
|   | Pemetrexed 500 mg powder for inj. SDV Ψ  |              |         |               | 68001-0544-xx |
| Pemetrexed 750 mg powder for inj. SDV Ψ                     | 68001-0545-xx                            |              |         |               |               |
| Pemetrexed 1000 mg powder for inj. SDV Ψ                    | 68001-0546-xx                            |              |         |               |               |
| Pemetrexed Disodium Solution for injection                  | Pemetrexed 100 mg/4 mL inj. SDV Ψ        | Sandoz       | Brand   | J9297         | 00781-3518-xx |
|   |  | Accord       | Brand   | J9296         | 16729-0522-xx |
|   |  | Hospira      | Brand   | J9294         | 00409-1045-xx |
|   | Pemetrexed 500 mg/20 mL inj. SDV Ψ       | Sandoz       | Brand   | J9297         | 00781-3519-xx |
|   |  | Accord       | Brand   | J9296         | 16729-0522-xx |
|   |  | Hospira      | Brand   | J9294         | 00409-2188-xx |
|   | Pemetrexed 850 mg/34mL inj. SDV Ψ        | Accord       | Brand   | J9296         | 16729-0522-xx |
|   | Pemetrexed 1000 mg/40 mL inj. SDV Ψ      | Accord       | Brand   | J9296         | 16729-0522-xx |
| Hospira   |  | Brand        | J9294   | 00409-3532-xx |               |
| Pemetrexed Solution for injection                           | Pemfexy 500 mg/20 mL inj. MDV            | Eagle        | Brand   | J9304         | 42367-0531-xx |
|   | Pemetrexed 100 mg/4mL inj. SDV Ψ         | Teva         | Brand   | J9314         | 00480-4516-xx |
|   | Pemetrexed 500 mg/20 mL inj. SDV Ψ       | Teva         | Brand   | J9314         | 00480-4514-xx |
|   | Pemetrexed 1000 mg/40 mL inj. SDV Ψ      | Teva         | Brand   | J9314         | 00480-4515-xx |
|   | Pemetrexed 100 mg powder for inj. SDV Ψ  | Hospira      | Brand   | J9323         | 00409-1060-xx |

|  |   |  |  |  |               |
|--|---|--|--|--|---------------|
| Pemetrexed<br>Ditromethamine for<br>injection  | Pemetrexed 500 mg powder for inj. SDV Ψ |  |  |  | 00409-1061-xx |
| <p><b>*Multiple manufacturers produce ANDA generics</b></p> <p>Ψ Designated products approved by the FDA as a 505(b)(2) NDA of the innovator product. These products are not rated as therapeutically equivalent to their reference listed drug in the Food and Drug Administration's (FDA) Orange Book and are therefore considered single source products based on the statutory definition of "single source drug" in section 1847A(c)(6) of the Act. For a complete list of all approved 505(b)(2) NDA products please reference the latest edition of the Orange Book: <a href="#">Approved Drug Products with Therapeutic Equivalence Evaluations   Orange Book   FDA</a></p>  |   |  |  |  |               |
| <p><b>J9294 – Injection, pemetrexed (hospira), not therapeutically equivalent to J9305, 10 mg</b><br/> <b>J9296 – Injection, pemetrexed (accord), not therapeutically equivalent to J9305, 10 mg</b><br/> <b>J9297 – Injection, pemetrexed (sandoz), not therapeutically equivalent to J9305, 10 mg</b><br/> <b>J9304 – Injection, pemetrexed (pemfexy), 10 mg</b><br/> <b>J9305 – Injection, pemetrexed, not otherwise specified, 10 mg</b><br/> <b>J9314 – Injection, pemetrexed (teva), not therapeutically equivalent to J9305, 10 mg</b><br/> <b>J9322 – Injection, pemetrexed (bluepoint), not therapeutically equivalent to J9305, 10 mg</b><br/> <b>J9323 – Injection, pemetrexed ditromethamine, 10 mg</b><br/> <b>J9324 – Injection, pemetrexed (pemrydi rtu), 10 mg</b><br/> <b>J9999 – Injection, pemetrexed various (shipla, etc.), 10 mg</b></p> |   |  |  |  |               |

## VII. References (STANDARD)

1. Alimta [package insert]. Indianapolis, IN; Eli Lilly; May 2023. Accessed June 2024.
2. Pemfexy [package insert]. Woodcliff Lake, NJ; Eagle Pharmaceuticals, Inc; December 2022. Accessed June 2024.
3. Pemrydi RTU [package insert]. Bridgewater, NJ; Amneal Pharmaceuticals LLC; June 2023. Accessed June 2024.
4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for pemetrexed. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.
5. Castagneto B, Botta M, Aitini E, et al, "Phase II Study of Pemetrexed in Combination With Carboplatin in Patients With Malignant Pleural Mesothelioma (MPM)," Ann Oncol, 2008, 19(2):370-3.
6. Ceresoli GL, Zucali PA, Favaretto AG, et al, "Phase II Study of Pemetrexed plus Carboplatin in Malignant Pleural Mesothelioma," J Clin Oncol, 2006, 24(9):1443-8.
7. Taylor P, Castagneto B, Dark G, et al, "Single-Agent Pemetrexed for Chemo-naïve and Pretreated Patients With Malignant Pleural Mesothelioma: Results of an International Expanded Access Program," J Thorac Oncol, 2008, 3(7):764-71.
8. Ciuleanu T, Brodowicz T, Zielinski C, et al, "Maintenance Pemetrexed Plus Best Supportive Care versus Placebo Plus Best Supportive Care for Non-Small-Cell Lung Cancer: A Randomised, Double-Blind, Phase 3 Study," Lancet, 2009, 374(9699):1432-40.
9. Grønberg BH, Bremnes RM, Fløtten O, et al, "Phase III Study by the Norwegian Lung Cancer Study Group: Pemetrexed Plus Carboplatin Compared With Gemcitabine Plus Carboplatin as

- First-Line Chemotherapy in Advanced Non-Small-Cell Lung Cancer,” J Clin Oncol, 2009, 27(19):3217-24.
10. Hanna N, Shepherd FA, Fossella FV, et al, “Randomized Phase III Trial of Pemetrexed versus Docetaxel in Patients With Non-Small-Cell Lung Cancer Previously Treated With Chemotherapy,” J Clin Oncol, 2004, 22(9):1589-97.
  11. Jassem J, Ramlau R, Santoro A, et al, “Phase III Trial of Pemetrexed Plus Best Supportive Care Compared With Best Supportive Care in Previously Treated Patients With Advanced Malignant Pleural Mesothelioma,” J Clin Oncol, 2008, 26(10):1698-704. [PubMed 18375898]
  12. Scagliotti GV, Parikh P, von Pawel J, et al, “Phase III Study Comparing Cisplatin Plus Gemcitabine With Cisplatin Plus Pemetrexed in Chemotherapy-Naive Patients With Advanced-Stage Non-Small-Cell Lung Cancer,” J Clin Oncol, 2008, 26(21):3543-51.
  13. Langer CJ, Gadgeel SM, Borghaei H, et al. Carboplatin and pemetrexed with or without pembrolizumab for advanced, non-squamous non-small-cell lung cancer: a randomised, phase 2 cohort of the open-label KEYNOTE-021 study. Lancet Oncol. 2016;17(11):1497-1508.
  14. Miller DS, Blessing JA, Krasner CN, et al, “Phase II Evaluation of Pemetrexed in the Treatment of Recurrent or Persistent Platinum-Resistant Ovarian or Primary Peritoneal Carcinoma: A Study of the Gynecologic Oncology Group,” J Clin Oncol, 2009, 27(16):2686-91.
  15. Liang Y, Padda SK, Riess JW, et al. Pemetrexed in patients with thymic malignancies previously treated with chemotherapy. Lung Cancer. 2015 Jan;87(1):34-8.
  16. Gbolahan OB, Porter RF, Salter JT, et al. A Phase II Study of Pemetrexed in Patients with Recurrent Thymoma and Thymic Carcinoma. J Thorac Oncol. 2018 Dec;13(12):1940-1948.
  17. Raizer JJ, Rademaker A, Evens AM, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. Cancer. 2012 Aug 1;118(15):3743-8.
  18. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. J Oncol Pract. 2018 Mar;14(3):e130-e136.
  19. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from [http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug\\_Waste\\_2019.pdf](http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf)
  20. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. BMJ. 2016 Feb 29;352:i788.
  21. Gandhi L, Rodríguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. N Engl J Med. 2018;378(22):2078-2092. doi:10.1056/NEJMoa1801005.
  22. Wu YL, Lu S, Cheng Y, et al. Efficacy and safety of pemetrexed/cisplatin versus gemcitabine/cisplatin as first-line treatment in Chinese patients with advanced nonsquamous non-small cell lung cancer. Lung Cancer. 2014;85(3):401-407. doi:10.1016/j.lungcan.2014.07.007.
  23. Paz-Ares L, de Marinis F, Dediu M, et al. Maintenance therapy with pemetrexed plus best supportive care versus placebo plus best supportive care after induction therapy with



- pemetrexed plus cisplatin for advanced non-squamous non-small-cell lung cancer (PARAMOUNT): a double-blind, phase 3, randomised controlled trial. *Lancet Oncol.* 2012;13(3):247-255. doi:10.1016/S1470-2045(12)70063-3.
24. Vogelzang NJ, Rusthoven JJ, Symanowski J, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol.* 2003;21(14):2636-2644. doi:10.1200/JCO.2003.11.136.
  25. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Ovarian Cancer/Fallopian Tube Cancer/Primary Peritoneal Cancer Version 2.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.
  26. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Thymomas and Thymic Carcinomas. Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.
  27. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Mesothelioma: Pleural Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.
  28. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Central Nervous System Cancers Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.
  29. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Non-Small Cell Lung Cancer Version 5.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2024.

30. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines<sup>®</sup>) for Mesothelioma: Peritoneal Version 1.2024. National Comprehensive Cancer Network, 2024. NATIONAL COMPREHENSIVE CANCER NETWORK<sup>®</sup>, NCCN<sup>®</sup>, and NCCN GUIDELINES<sup>®</sup> are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2024.
31. Forde P, Spicer J, Provencio M, et.al. Abstract CT003: Nivolumab (NIVO) + platinum-doublet chemotherapy (chemo) vs chemo as neoadjuvant treatment (tx) for resectable (IB-IIIa) non-small cell lung cancer (NSCLC) in the phase 3 CheckMate 816 trial. *Cancer Res* (2021) 81 (13\_Supplement): CT003. <https://doi.org/10.1158/1538-7445.AM2021-CT003>
32. Miller DS, Blessing JA, Bodurka DC, et al. Evaluation of pemetrexed (Alimta, LY231514) as second line chemotherapy in persistent or recurrent carcinoma of the cervix: a phase II study of the Gynecologic Oncology Group. *Gynecol Oncol*. 2008 Jul;110(1):65-70. doi: 10.1016/j.ygyno.2008.03.009.
33. Zalcman G, Mazieres J, Margery J, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2016 Apr 2;387(10026):1405-1414. doi: 10.1016/S0140-6736(15)01238-6.
34. Fan C, Zhao Q, Li L, et al. Efficacy and Safety of Intrathecal Pemetrexed Combined With Dexamethasone for Treating Tyrosine Kinase Inhibitor-Failed Leptomeningeal Metastases From EGFR-Mutant NSCLC-a Prospective, Open-Label, Single-Arm Phase 1/2 Clinical Trial (Unique Identifier: ChiCTR1800016615). *J Thorac Oncol*. 2021 Aug;16(8):1359-1368. doi: 10.1016/j.jtho.2021.04.018.
35. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines<sup>®</sup>) for Cervical Cancer Version 3.2024. National Comprehensive Cancer Network, 2024. NATIONAL COMPREHENSIVE CANCER NETWORK<sup>®</sup>, NCCN<sup>®</sup>, and NCCN GUIDELINES<sup>®</sup> are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2024.
36. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines<sup>®</sup>) for Vaginal Cancer Version 1.2025. National Comprehensive Cancer Network, 2024. NATIONAL COMPREHENSIVE CANCER NETWORK<sup>®</sup>, NCCN<sup>®</sup>, and NCCN GUIDELINES<sup>®</sup> are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2024.

## VIII. References (ENHANCED)

- 1e. Sweeney CJ, Roth BJ, Kabbinavar FF, et al. Phase II study of pemetrexed for second-line treatment of transitional cell cancer of the urothelium. *J Clin Oncol*. 2006 Jul 20;24(21):3451-7.
- 2e. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med*. 2017;376(11):1015–1026.

- 3e. Rosenberg JE, Hoffman-Censits J, Powles T, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet*. 2016;387(10031):1909–1920.
- 4e. Powles T, Durán I, van der Heijden MS, et al. Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2018 Feb 24;391(10122):748-757.
- 5e. Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2017 Mar;18(3):312-322.
- 6e. Massard C, Gordon MS, Sharma S, et al. Safety and Efficacy of Durvalumab (MEDI4736), an Anti-Programmed Cell Death Ligand-1 Immune Checkpoint Inhibitor, in Patients With Advanced Urothelial Bladder Cancer. *J Clin Oncol*. 2016;34(26):3119–3125.
- 7e. Patel MR, Ellerton J, Infante J, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol*. 2018 Jan;19(1):51-64.
- 8e. Ko YJ, et al. Nanoparticle albumin-bound paclitaxel for second-line treatment of metastatic urothelial carcinoma: a single group, multicentre, phase 2 study. *Lancet Oncol*. 2013 Jul;14(8):769-76.
- 9e. Lorusso V, et al. A phase II study of gemcitabine in patients with transitional cell carcinoma of the urinary tract previously treated with platinum. Italian Co-operative Group on Bladder Cancer. *Eur J Cancer*. 1998 Jul;34(8):1208-12.
- 10e. Meluch AA, et al. Paclitaxel and gemcitabine chemotherapy for advanced transitional-cell carcinoma of the urothelial tract: a phase II trial of the Minnie pearl cancer research network.. *J Clin Oncol*. 2001 Jun 15;19(12):3018-24.
- 11e. von der Maase H, et al. Gemcitabine and Cisplatin Versus Methotrexate, Vinblastine, Doxorubicin, and Cisplatin in Advanced or Metastatic Bladder Cancer: Results of a Large, Randomized, Multinational, Multicenter, Phase III Study. *Journal of Clinical Oncology* 2000 18:17, 3068-3077.
- 12e. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/ carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer "unfit" for cisplatin-based chemotherapy: phase II--results of EORTC study 30986. *J Clin Oncol*. 2009;27(33):5634–5639.
- 13e. McCaffrey JA, et al. Phase II trial of docetaxel in patients with advanced or metastatic transitional-cell carcinoma. *J Clin Oncol*. 1997 May;15(5):1853-7.
- 14e. Vaughn DJ, et al. Phase II trial of weekly paclitaxel in patients with previously treated advanced urothelial cancer. *J Clin Oncol*. 2002 Feb 15;20(4):937-40.

- 15e. Petrylak DP, et al. Ramucirumab plus docetaxel versus placebo plus docetaxel in patients with locally advanced or metastatic urothelial carcinoma after platinum-based therapy (RANGE): a randomised, double-blind, phase 3 trial. *Lancet*. 2017 Nov 18;390(10109):2266-2277.
- 16e. Witte RS, et al. Eastern Cooperative Oncology Group phase II trial of ifosfamide in the treatment of previously treated advanced urothelial carcinoma. *J Clin Oncol*. 1997 Feb;15(2):589-93.
- 17e. Siefker-Radtke AO, Dinney CP, Shen Y, et al. A phase 2 clinical trial of sequential neoadjuvant chemotherapy with ifosfamide, doxorubicin, and gemcitabine followed by cisplatin, gemcitabine, and ifosfamide in locally advanced urothelial cancer: final results. *Cancer*. 2012;119(3):540-7.
- 18e. Sternberg CN, et al. Randomized phase III trial of high-dose-intensity methotrexate, vinblastine, doxorubicin, and cisplatin (MVAC) chemotherapy and recombinant human granulocyte colony-stimulating factor versus classic MVAC in advanced urothelial tract tumors: European Organization for Research and Treatment of Cancer Protocol no. 30924. *J Clin Oncol*. 2001 May 15;19(10):2638-46.
- 19e. Plotkin SR, Betensky RA, Hochberg FH, et al. Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. *Clin Cancer Res*. 2004 Sep 1;10(17):5643-6.
- 20e. Nayak L, Abrey LE, Drappatz J, et al. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma*. 2013;54(1):58–61.
- 21e. Krug LM, Pass HI, Rusch VW, et al. Multicenter phase II trial of neoadjuvant pemetrexed plus cisplatin followed by extrapleural pneumonectomy and radiation for malignant pleural mesothelioma. *J Clin Oncol*. 2009;27(18):3007–3013.
- 22e. Santoro A, O'Brien ME, Stahel RA, et al. Pemetrexed plus cisplatin or pemetrexed plus carboplatin for chemo-naïve patients with malignant pleural mesothelioma: results of the International Expanded Access Program. *J Thorac Oncol*. 2008 Jul;3(7):756-63.
- 23e. Zalcman G, Mazieres J, Margery J, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2016 Apr 2;387(10026):1405-1414.
- 24e. Ceresoli GL, Zucali PA, Mencoboni M, et al. Phase II study of pemetrexed and carboplatin plus bevacizumab as first-line therapy in malignant pleural mesothelioma. *Br J Cancer*. 2013;109(3):552–558.
- 25e. Muers MF, Stephens RJ, Fisher P, et al. Active symptom control with or without chemotherapy in the treatment of patients with malignant pleural mesothelioma (MS01): a multicentre randomised trial. *Lancet*. 2008;371(9625):1685–1694.
- 26e. Zucali PA, Simonelli M, Michetti G, et al. Second-line chemotherapy in malignant pleural mesothelioma: results of a retrospective multicenter survey. *Lung Cancer*. 2012 Mar;75(3):360-7.
- 27e. Scherpereel A, Mazieres J, Greillier L, et al. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. *Lancet Oncol*. 2019 Feb;20(2):239-253.

- 28e. Scherpereel A, Mazieres J, Greillier L, et al. Second or 3rd line nivolumab (Nivo) versus nivo plus ipilimumab (Ipi) in malignant pleural mesothelioma (MPM) patients: Updated results of the IFCT-1501 MAPS2 randomized phase 2 trial. *Ann Oncol*. 2017 Sept;28(5):mdx440.074.
- 29e. Disselhorst MJ, Quispel-Janssen J, Lalezari F, et al. Ipilimumab and nivolumab in the treatment of recurrent malignant pleural mesothelioma (INITIATE): results of a prospective, single-arm, phase 2 trial. *Lancet Respir Med*. 2019 Mar;7(3):260-270.
- 30e. Quispel-Janssen J, van der Noort V, de Vries JF, et al. Programmed Death 1 Blockade With Nivolumab in Patients With Recurrent Malignant Pleural Mesothelioma. *J Thorac Oncol*. 2018 Oct;13(10):1569-1576.
- 31e. Alley EW, Lopez J, Santoro A, et al. Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial. *Lancet Oncol*. 2017 May;18(5):623-630.
- 32e. Alley EW, Lopez J, Santoro A, et al. Long-Term Overall Survival for Patients with Malignant Pleural Mesothelioma on Pembrolizumab Enrolled in KEYNOTE-028. *J Thorac Oncol*. 2017 Jan;12(1):S294.
- 33e. Metaxas Y, Rivalland G, Mauti LA, et al. Pembrolizumab as Palliative Immunotherapy in Malignant Pleural Mesothelioma. *J Thorac Oncol*. 2018 Nov;13(11):1784-1791.
- 34e. Stebbing J, Powles T, McPherson K, et al. The efficacy and safety of weekly vinorelbine in relapsed malignant pleural mesothelioma. *Lung Cancer*. 2009 Jan;63(1):94-7.
- 35e. Zauderer MG, Kass SL, Woo K, Sima CS, Ginsberg MS, Krug LM. Vinorelbine and gemcitabine as second- or third-line therapy for malignant pleural mesothelioma. *Lung Cancer*. 2014;84(3):271–274.
- 36e. Kim JS, Lim SY, Hwang J, Kang EJ, Choi YJ. A Case Report of Primary Pericardial Malignant Mesothelioma Treated with Pemetrexed and Cisplatin. *J Korean Med Sci*. 2017;32(11):1879–1884.
- 37e. Carteni G, Manegold C, Garcia GM, et al. Malignant peritoneal mesothelioma-Results from the International Expanded Access Program using pemetrexed alone or in combination with a platinum agent. *Lung Cancer*. 2009 May;64(2):211-8.
- 38e. Zhang L, Ou W, Liu Q, Li N, Liu L, Wang S. Pemetrexed plus carboplatin as adjuvant chemotherapy in patients with curative resected non-squamous non-small cell lung cancer. *Thorac Cancer*. 2014;5(1):50–56.
- 39e. Kreuter M, Vansteenkiste J, Fischer JR, et al. Randomized phase 2 trial on refinement of early-stage NSCLC adjuvant chemotherapy with cisplatin and pemetrexed versus cisplatin and vinorelbine: the TREAT study. *Ann Oncol*. 2013 Apr;24(4):986-92.
- 40e. Kenmotsu H, Yamamoto N, Yamanaka T, et al. Randomized phase III study of pemetrexed/cisplatin (Pem/Cis) versus vinorelbine /cisplatin (Vnr/Cis) for completely resected stage II-IIIa non-squamous non-small-cell lung cancer (Ns-NSCLC): The JIPANG study. *J Clin Oncol*, 2019; 37(15\_suppl):8501.
- 41e. Arriagada R, Bergman B, Dunant A, et al. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *N Engl J Med*. 2004 Jan 22;350(4):351-60.

- 42e. Scagliotti GV, Pastorino U, Vansteenkiste JF, et al. Randomized phase III study of surgery alone or surgery plus preoperative cisplatin and gemcitabine in stages IB to IIIA non-small-cell lung cancer. *J Clin Oncol*. 2012 Jan 10;30(2):172-8.
- 43e. Strauss GM, Herndon JE 2nd, Maddaus MA, et al. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small-cell lung cancer: CALGB 9633 with the Cancer and Leukemia Group B, Radiation Therapy Oncology Group, and North Central Cancer Treatment Group Study Groups. *J Clin Oncol*. 2008;26(31):5043–5051.
- 44e. Usami N, Yokoi K, Hasegawa Y, et al. Phase II study of carboplatin and gemcitabine as adjuvant chemotherapy in patients with completely resected non-small cell lung cancer: a report from the Central Japan Lung Study Group, CJLSG 0503 trial. *Int J Clin Oncol*. 2010 Dec;15(6):583-7.
- 45e. Senan S, Brade A, Wang LH, et al. PROCLAIM: Randomized Phase III Trial of Pemetrexed-Cisplatin or Etoposide-Cisplatin Plus Thoracic Radiation Therapy Followed by Consolidation Chemotherapy in Locally Advanced Nonsquamous Non-Small-Cell Lung Cancer. *J Clin Oncol*. 2016 Mar 20;34(9):953-62.
- 46e. Curran WJ Jr, Paulus R, Langer CJ, et al. Sequential vs. concurrent chemoradiation for stage III non-small cell lung cancer: randomized phase III trial RTOG 9410 [published correction appears in *J Natl Cancer Inst*. 2012 Jan 4;104(1):79]. *J Natl Cancer Inst*. 2011;103(19):1452–1460.
- 47e. Belani CP, Choy H, Bonomi P, et al. Combined chemoradiotherapy regimens of paclitaxel and carboplatin for locally advanced non-small-cell lung cancer: a randomized phase II locally advanced multi-modality protocol. *J Clin Oncol*. 2005 Sep 1;23(25):5883-91.
- 48e. Yang JC, Hirsh V, Schuler M, et al. Symptom control and quality of life in LUX-Lung 3: a phase III study of afatinib or cisplatin/pemetrexed in patients with advanced lung adenocarcinoma with EGFR mutations. *J Clin Oncol*. 2013 Sep 20;31(27):3342-50.
- 49e. Zukin M, Barrios CH, Pereira JR, et al. Randomized phase III trial of single-agent pemetrexed versus carboplatin and pemetrexed in patients with advanced non-small-cell lung cancer and Eastern Cooperative Oncology Group performance status of 2. *J Clin Oncol*. 2013 Aug 10;31(23):2849-53.
- 50e. Gridelli C, Kaukel E, Gregorc V, et al. Single-agent pemetrexed or sequential pemetrexed/gemcitabine as front-line treatment of advanced non-small cell lung cancer in elderly patients or patients ineligible for platinum-based chemotherapy: a multicenter, randomized, phase II trial. *J Thorac Oncol*. 2007 Mar;2(3):221-9.
- 51e. Rusthoven JJ, Eisenhauer E, Butts C, et al. Multitargeted antifolate LY231514 as first-line chemotherapy for patients with advanced non-small-cell lung cancer: A phase II study. National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol*. 1999 Apr;17(4):1194.
- 52e. Patel JD, Socinski MA, Garon EB, et al. PointBreak: a randomized phase III study of pemetrexed plus carboplatin and bevacizumab followed by maintenance pemetrexed and bevacizumab versus paclitaxel plus carboplatin and bevacizumab followed by maintenance

bevacizumab in patients with stage IIIB or IV nonsquamous non-small-cell lung cancer. *J Clin Oncol*. 2013;31(34):4349–4357.

- 53e. Barlesi F, Scherpereel A, Rittmeyer A, et al. Randomized phase III trial of maintenance bevacizumab with or without pemetrexed after first-line induction with bevacizumab, cisplatin, and pemetrexed in advanced nonsquamous non-small-cell lung cancer: AVAPERL (MO22089). *J Clin Oncol*. 2013 Aug 20;31(24):3004-11.
- 54e. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1–Positive Non–Small-Cell Lung Cancer. *N Engl J Med* 2016; 375:1823-1833.
- 55e. Socinski MA, Jotte RM, Capuzzo F, et al. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018; 378:2288-2301.
- 56e. Cardenal F, López-Cabrerizo MP, Antón A, et al. Randomized phase III study of gemcitabine-cisplatin versus etoposide-cisplatin in the treatment of locally advanced or metastatic non-small-cell lung cancer. *J Clin Oncol*. 1999 Jan;17(1):12-8.
- 57e. Fossella F, Pereira JR, von Pawel J, et al. Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. *J Clin Oncol*. 2003 Aug 15;21(16):3016-24.
- 58e. Zatloukal P, Kanitz E, Magyar P, et al. Gemcitabine in locally advanced and metastatic non-small cell lung cancer: the Central European phase II study. *Lung Cancer*. 1998 Dec;22(3):243-50.
- 59e. Pujol JL, Breton JL, Gervais R, et al. Gemcitabine-docetaxel versus cisplatin-vinorelbine in advanced or metastatic non-small-cell lung cancer: a phase III study addressing the case for cisplatin. *Ann Oncol*. 2005 Apr;16(4):602-10.
- 60e. Tan EH, Szczesna A, Krzakowski M, et al. Randomized study of vinorelbine--gemcitabine versus vinorelbine--carboplatin in patients with advanced non-small cell lung cancer. *Lung Cancer*. 2005 Aug;49(2):233-40.
- 61e. Paz-Ares L, de Marinis F, Dediu M, et al. PARAMOUNT: Final Overall Survival Results of the Phase III Study of Maintenance Pemetrexed Versus Placebo Immediately After Induction Treatment With Pemetrexed Plus Cisplatin for Advanced Nonsquamous Non–Small-Cell Lung Cancer. *J Clin Oncol*. 2013 Aug 10;31(23):2895-902.
- 62e. Anderson H, Hopwood P, Stephens RJ, et al. Gemcitabine plus best supportive care (BSC) vs BSC in inoperable non-small cell lung cancer--a randomized trial with quality of life as the primary outcome. UK NSCLC Gemcitabine Group. *Non-Small Cell Lung Cancer*. *Br J Cancer*. 2000;83(4):447–453.
- 63e. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med*. 2015;373(17):1627–1639.
- 64e. Herbst RS, Baas P, Kim DW, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet*. 2016 Apr 9;387(10027):1540-50.

- 65e. Barlesi F, Park K, Ciardiello F, et al. Primary analysis from OAK, a randomized phase III study comparing atezolizumab with docetaxel in 2L/3L NSCLC. *Ann Oncol*. 2016 Oct;27(6):LBA44\_PR.
- 66e. Garon EB, Ciuleanu TE, Arrieta O, et al. Ramucirumab plus docetaxel versus placebo plus docetaxel for second-line treatment of stage IV non-small-cell lung cancer after disease progression on platinum-based therapy (REVEL): a multicentre, double-blind, randomised phase 3 trial. *Lancet*. 2014 Aug 23;384(9944):665-73.
- 67e. Ceresoli GL, Gregorc V, Cordio S, et al. Phase II study of weekly paclitaxel as second-line therapy in patients with advanced non-small cell lung cancer. *Lung Cancer*. 2004 May;44(2):231-9.
- 68e. Palmieri G, Buonerba C, Ottaviano M, et al. Capecitabine plus gemcitabine in thymic epithelial tumors: final analysis of a Phase II trial. *Future Oncol*. 2014 Nov;10(14):2141-7.
- 69e. Thomas A, Rajan A, Berman A, et al. Sunitinib in patients with chemotherapy-refractory thymoma and thymic carcinoma: an open-label phase 2 trial [published correction appears in *Lancet Oncol*. 2015 Mar;16(3):e105]. *Lancet Oncol*. 2015;16(2):177–186.
- 70e. Zucali PA, De Pas T, Palmieri G, et al. Phase II Study of Everolimus in Patients With Thymoma and Thymic Carcinoma Previously Treated With Cisplatin-Based Chemotherapy. *J Clin Oncol*. 2018 Feb 1;36(4):342-349.
- 71e. Loehrer PJ Sr, Wang W, Johnson DH, et al. Octreotide alone or with prednisone in patients with advanced thymoma and thymic carcinoma: an Eastern Cooperative Oncology Group Phase II Trial. *J Clin Oncol*. 2004 Jan 15;22(2):293-9.
- 72e. Umemura S, Segawa Y, Fujiwara K, et al. A case of recurrent metastatic thymoma showing a marked response to paclitaxel monotherapy. *Jpn J Clin Oncol*. 2002 Jul;32(7):262-5.
- 73e. Bluthgen MV, Boutros C, Fayard F, et al. Activity and safety of oral etoposide in pretreated patients with metastatic or recurrent thymic epithelial tumors (TET): A single-institution experience. *Lung Cancer*. 2016 Sep;99:111-6.
- 74e. Rose PG, Blessing JA, Ball HG, et al. A phase II study of docetaxel in paclitaxel-resistant ovarian and peritoneal carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol*. 2003 Feb;88(2):130-5.
- 75e. Rose PG, Blessing JA, Mayer AR, Homesley HD. Prolonged oral etoposide as second-line therapy for platinum-resistant and platinum-sensitive ovarian carcinoma: a Gynecologic Oncology Group study. *J Clin Oncol*. 1998 Feb;16(2):405-10.
- 76e. Gordon AN, Tonda M, Sun S, et al. Long-term survival advantage for women treated with pegylated liposomal doxorubicin compared with topotecan in a phase 3 randomized study of recurrent and refractory epithelial ovarian cancer. *Gynecol Oncol*. 2004 Oct;95(1):1-8.
- 77e. Sehouli J, Stengel D, Harter P, et al. Topotecan Weekly Versus Conventional 5-Day Schedule in Patients With Platinum-Resistant Ovarian Cancer: a randomized multicenter phase II trial of the North-Eastern German Society of Gynecological Oncology Ovarian Cancer Study Group. *J Clin Oncol*. 2011 Jan 10;29(2):242-8.



- 78e. Kindler, HL, Ismaila N, Armato III, SG, et al. Treatment of Malignant Pleural Mesothelioma: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol*. 2018 Jan;36(13):1343-1373.
- 79e. Nowak AK, Byrne MJ, Williamson R, et al. A multicentre phase II study of cisplatin and gemcitabine for malignant mesothelioma. *Br J Cancer*. 2002;87(5):491-496.
- 80e. van Haarst JM, Baas P, Manegold Ch, et al. Multicentre phase II study of gemcitabine and cisplatin in malignant pleural mesothelioma. *Br J Cancer*. 2002;86(3):342-345.
- 81e. Spigel D et al. IMpower110: Interim OS Analysis of a Phase III Study of Atezolizumab (atezo) vs Platinum-Based Chemotherapy (chemo) as 1L Treatment (tx) in PD-L1–selected NSCLC [ESMO 2019 Abstract LBA78].
- 82e. Reck M, Ciuleanu T-E, Dols MC, et al. Nivolumab (NIVO) + ipilimumab (IPI) + 2 cycles of platinum-doublet chemotherapy (chemo) vs 4 cycles chemo as first-line (1L) treatment (tx) for stage IV/recurrent non-small cell lung cancer (NSCLC): CheckMate 9LA [abstract]. *J Clin Oncol* 2020;38:Abstract 9501-9501.
- 83e. West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*. 2019;20(7):924-937.
- 84e. Zalcman G, Peters S, Mansfield AS, et al. Checkmate 743: A phase 3, randomized, open-label trial of nivolumab (nivo) plus ipilimumab (ipi) vs pemetrexed plus cisplatin or carboplatin as first-line therapy in unresectable pleural mesothelioma. *Journal of Clinical Oncology* 2017 35:15\_suppl, TPS8581-TPS8581.
- 85e. Sezer A, Kilickap S, Gümüş M, et al. Cemiplimab monotherapy for first-line treatment of advanced non-small-cell lung cancer with PD-L1 of at least 50%: a multicentre, open-label, global, phase 3, randomised, controlled trial. *Lancet*. 2021 Feb 13;397(10274):592-604.
- 86e. Scagliotti GV, Shin DM, Kindler HL, et al. Phase II study of pemetrexed with and without folic acid and vitamin B12 as front-line therapy in malignant pleural mesothelioma. *J Clin Oncol*. 2003 Apr 15;21(8):1556-61.
- 87e. Jänne PA, Wozniak AJ, Belani CP, et al. Open-label study of pemetrexed alone or in combination with cisplatin for the treatment of patients with peritoneal mesothelioma: outcomes of an expanded access program. *Clin Lung Cancer*. 2005 Jul;7(1):40-6.
- 88e. Gogishvili M, Melkadze T, Makharadze T, et al. LBA51 EMPOWER-Lung 3: Cemiplimab in combination with platinum doublet chemotherapy for first-line (1L) treatment of advanced non-small cell lung cancer (NSCLC). *Annals of Oncology*, ISSN: 0923-7534, Vol: 32, SUPPLEMENT 5, S1328, SEPTEMBER 01, 2021. DOI10.1016/j.annonc.2021.08.2130.
- 89e. Johnson ML, Cho BC, Luft A, et al. Durvalumab With or Without Tremelimumab in Combination With Chemotherapy as First-Line Therapy for Metastatic Non-Small-Cell Lung Cancer: The Phase III POSEIDON Study. *J Clin Oncol*. 2023 Feb 20;41(6):1213-1227.

- 90e. Bijelic L, Stuart OA, Sugarbaker P. Adjuvant bidirectional chemotherapy with intraperitoneal pemetrexed combined with intravenous Cisplatin for diffuse malignant peritoneal mesothelioma. *Gastroenterol Res Pract*. 2012;2012:890450.
- 91e. Wakelee H, Liberman M, Kato T, et al. Perioperative Pembrolizumab for Early-Stage Non-Small-Cell Lung Cancer. *N Engl J Med*. 2023 Aug 10;389(6):491-503. doi: 10.1056/NEJMoa2302983. Epub 2023 Jun 3. PMID: 37272513.
- 92e. Novello S, Mazières J, Oh IJ, et al. Alectinib versus chemotherapy in crizotinib-pretreated anaplastic lymphoma kinase (ALK)-positive non-small-cell lung cancer: results from the phase III ALUR study. *Ann Oncol*. 2018;29(6):1409-1416. doi:10.1093/annonc/mdy121.
- 93e. Kim DW, Tiseo M, Ahn MJ, et al. Brigatinib in Patients With Crizotinib-Refractory Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer: A Randomized, Multicenter Phase II Trial. *J Clin Oncol*. 2017 Aug 1;35(22):2490-2498. doi: 10.1200/JCO.2016.71.5904.
- 94e. Shaw AT, Kim DW, Nakagawa K, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med*. 2013 Jun 20;368(25):2385-94. doi: 10.1056/NEJMoa1214886.
- 95e. Solomon BJ, Besse B, Bauer TM, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study [published correction appears in *Lancet Oncol*. 2019 Jan;20(1):e10]. *Lancet Oncol*. 2018;19(12):1654–1667. doi:10.1016/S1470-2045(18)30649-1.
- 96e. Planchard D, Jänne PA, Cheng Y, et al; FLAURA2 Investigators. Osimertinib with or without Chemotherapy in EGFR-Mutated Advanced NSCLC. *N Engl J Med*. 2023 Nov 23;389(21):1935-1948. doi: 10.1056/NEJMoa2306434. Epub 2023 Nov 8.
- 97e. Zhou C, Tang KJ, Cho BC, et al; PAPILLON Investigators. Amivantamab plus Chemotherapy in NSCLC with EGFR Exon 20 Insertions. *N Engl J Med*. 2023 Nov 30;389(22):2039-2051. doi: 10.1056/NEJMoa2306441. Epub 2023 Oct 21.
- 98e. Passaro A, Wang J, Wang Y, et al. Amivantamab plus chemotherapy with and without lazertinib in EGFR-mutant advanced NSCLC after disease progression on osimertinib: Primary results from the phase 3 MARIPOSA-2 study. *Annals of Oncology*. Published online October 1, 2023. doi:https://doi.org/10.1016/j.annonc.2023.10.117.
- 99e. Hainsworth JD, Waterhouse DM, Shih KC, et al. Phase II trial of preoperative pemetrexed plus carboplatin in patients with stage IB-III nonsquamous non-small cell lung cancer (NSCLC). *Lung Cancer*. 2018;118:6-12. doi:10.1016/j.lungcan.2018.01.009.
- 100e. Prime Therapeutics Management. Pemetrexed Clinical Literature Review Analysis. Last updated June 2024. Accessed June 2024.

## Appendix 1 – Covered Diagnosis Codes

| ICD-10 | ICD-10 Description                              |
|--------|---|
| C33    | Malignant neoplasm of trachea                   |
| C34.00 | Malignant neoplasm of unspecified main bronchus |

| ICD-10 | ICD-10 Description  |
|--------|---|
| C34.01 | Malignant neoplasm of right main bronchus                                 |
| C34.02 | Malignant neoplasm of left main bronchus                                  |
| C34.10 | Malignant neoplasm of upper lobe, unspecified bronchus or lung            |
| C34.11 | Malignant neoplasm of upper lobe, right bronchus or lung                  |
| C34.12 | Malignant neoplasm of upper lobe, left bronchus or lung                   |
| C34.2  | Malignant neoplasm of middle lobe, bronchus or lung                       |
| C34.30 | Malignant neoplasm of lower lobe, unspecified bronchus or lung            |
| C34.31 | Malignant neoplasm of lower lobe, right bronchus or lung                  |
| C34.32 | Malignant neoplasm of lower lobe, left bronchus or lung                   |
| C34.80 | Malignant neoplasm of overlapping sites of unspecified bronchus or lung   |
| C34.81 | Malignant neoplasm of overlapping sites of right bronchus and lung        |
| C34.82 | Malignant neoplasm of overlapping sites of left bronchus and lung         |
| C34.90 | Malignant neoplasm of unspecified part of unspecified bronchus or lung    |
| C34.91 | Malignant neoplasm of unspecified part of right bronchus or lung          |
| C34.92 | Malignant neoplasm of unspecified part of left bronchus or lung           |
| C37    | Malignant neoplasm of thymus  |
| C45.0  | Mesothelioma of pleura  |
| C45.1  | Mesothelioma of peritoneum  |
| C48.1  | Malignant neoplasm of specified parts of peritoneum                       |
| C48.2  | Malignant neoplasm of peritoneum, unspecified                             |
| C48.8  | Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum |
| C56.1  | Malignant neoplasm of right ovary   |
| C56.2  | Malignant neoplasm of left ovary  |
| C56.3  | Malignant neoplasm of bilateral ovaries                                   |
| C56.9  | Malignant neoplasm of unspecified ovary                                   |
| C57.00 | Malignant neoplasm of unspecified fallopian tube                          |
| C57.01 | Malignant neoplasm of right fallopian tube                                |
| C57.02 | Malignant neoplasm of left fallopian tube                                 |
| C57.10 | Malignant neoplasm of unspecified broad ligament                          |
| C57.11 | Malignant neoplasm of right broad ligament                                |
| C57.12 | Malignant neoplasm of left broad ligament                                 |
| C57.20 | Malignant neoplasm of unspecified round ligament                          |
| C57.21 | Malignant neoplasm of right round ligament                                |
| C57.22 | Malignant neoplasm of left round ligament                                 |

| ICD-10  | ICD-10 Description  |
|---------|---|
| C57.3   | Malignant neoplasm of parametrium                                 |
| C57.4   | Malignant neoplasm of uterine adnexa, unspecified                 |
| C57.7   | Malignant neoplasm of other specified female genital organs       |
| C57.8   | Malignant neoplasm of overlapping sites of female genital organs  |
| C57.9   | Malignant neoplasm of female genital organ, unspecified           |
| D15.0   | Benign neoplasm of thymus   |
| D38.4   | Neoplasm of uncertain behavior of thymus                          |
| Z85.118 | Personal history of other malignant neoplasm of bronchus and lung |
| Z85.238 | Personal history of other malignant neoplasm of thymus            |
| Z85.43  | Personal history of malignant neoplasm of ovary                   |

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

| Medicare Part B Administrative Contractor (MAC) Jurisdictions |   |   |
|---|---|---|
| Jurisdiction  | Applicable State/US Territory   | Contractor  |
| E (1)   | CA, HI, NV, AS, GU, CNMI  | Noridian Healthcare Solutions, LLC                |
| F (2 & 3)   | AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ  | Noridian Healthcare Solutions, LLC                |
| 5   | KS, NE, IA, MO  | Wisconsin Physicians Service Insurance Corp (WPS) |
| 6   | MN, WI, IL  | National Government Services, Inc. (NGS)          |
| H (4 & 7)   | LA, AR, MS, TX, OK, CO, NM  | Novitas Solutions, Inc.                           |
| 8   | MI, IN  | Wisconsin Physicians Service Insurance Corp (WPS) |
| N (9)   | FL, PR, VI  | First Coast Service Options, Inc.                 |
| J (10)  | TN, GA, AL  | Palmetto GBA                                      |
| M (11)  | NC, SC, WV, VA (excluding below)  | Palmetto GBA                                      |
| L (12)  | DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA) | Novitas Solutions, Inc.                           |
| K (13 & 14)   | NY, CT, MA, RI, VT, ME, NH  | National Government Services, Inc. (NGS)          |

| Medicare Part B Administrative Contractor (MAC) Jurisdictions |                               |                         |
|---|-------------------------------|-------------------------|
| Jurisdiction  | Applicable State/US Territory | Contractor              |
| 15  | KY, OH                        | CGS Administrators, LLC |